# PediaSure® Peptide 1.0 Cal

Peptide-Based Nutrition for Children

## **Nutrition Information - Liquid Vanilla**

	8 fl oz (237 mL)	
	Value	%RDI / %DV
Chromium, mcg	7.1	
Molybdenum, mcg	8.5	

## **Preparation**

#### Instructions for Use: 8-fl-oz Bottle

- Shake very well.
- 1 Best served chilled.
- Store unopened at room temperature.
- 1 After opening, refrigerate and use within 48 hours.

#### Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to child's condition and tolerance.
- Additional fluid may be required.

#### Oral Feeding:

1 Best when chilled.

#### Instructions for Use: Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and may reduce hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Follow directions for use provided by manufacturer of feeding sets.

### For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on feeding set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.

