

Product Information: Pivot® 1.5 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Pivot® 1.5 Cal

Peptide-based, high protein, therapeutic nutrition for metabolic stress

- PIVOT 1.5 CAL is peptide-based complete, balanced therapeutic nutrition for short- or long-term tube feeding for metabolically stressed surgical, trauma, burn, or head and neck cancer patients who could benefit from an immune-modulating enteral formula.
- For tube feeding.
- For sole-source nutrition.
- Use under medical supervision.
- Provides very high protein (93.8 g/L, 25% of calories) to support protein synthesis, tissue repair and wound healing.
- Provides 1.5 Cal/mL—concentrated calories for fluid-restricted patients.
- Immune support:
 - Arginine (13 g/L, 3.5% of calories) to support proliferation and function of immune cells.
 - Glutamine (inherent) (7.6 g/L) for GI-tract integrity and energy for immune cells.
 - Omega-3 fatty acids (EPA, 2.6 g/L; DHA, 1.1 g/L) to help modulate inflammation and support immune function.^{1,2}
- Tolerance:
 - Advanced blend of hydrolyzed protein, structured lipid, and prebiotic (NutraFlora® scFOS®) to promote absorption and tolerance.
 - Hydrolyzed, peptide-based protein system.
 - MCT/fish oil structured lipid, a well-tolerated^{3,4} and absorbed⁴ next generation fat to promote absorption of fatty acids.
 - 1.8 g of scFOS/8 fl oz (7.5 g/L). scFOS are prebiotic soluble fibers that stimulate the growth of beneficial bacteria in the colon.
- Elevated antioxidants vitamins C & E, and beta-carotene.
- Meets or exceeds 100% of the 1995 RDIs for 24 vitamins and minerals in 1500 Cal (1000 mL).
- Halal.
- Gluten-free.
- Suitable for lactose intolerance.



Safety Precautions

- Not for IV use.
- Not for people with galactosemia.

* NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.

¹ Calder PC. Prostaglandins Leukot Essent Fatty Acids. 2008;79:101-108.

² Calder PC. Clin Nutr. 2010;29:5-12.

³ Kenler AS, et al. Ann Surg. 1996;223:316-333.

⁴ McKenna MC, et al. J Pediatr Gastroenterol Nutr. 1985;4:45-51.

Ingredients

Liquid Unflavored:

Water, Corn Syrup Solids, Hydrolyzed Sodium Caseinate, Whey Protein Hydrolysate, Structured lipid (Interesterified Marine Oil [Contains One or More of the Following: Anchovy, Cod, Menhaden, Pollock, Salmon, Sardine, Tuna] and Medium Chain Triglycerides), Soy Oil, Canola Oil, L-Arginine, Fructooligosaccharides, Potassium Citrate, Citric Acid, Calcium Phosphate, Magnesium Chloride, Soy Lecithin, Natural & Artificial Flavor, Ascorbic Acid, Cellulose Gel, Choline Chloride, Magnesium Phosphate, Potassium Chloride, Carrageenan, Taurine, d-Alpha-Tocopheryl Acetate, L-Carnitine, Cellulose Gum, Zinc Sulfate, Ferrous Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Beta-Carotene, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin B12, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Pivot® 1.5 Cal

Peptide-based, high protein, therapeutic nutrition for metabolic stress

Availability

List Number	Item
62719	Pivot 1.5 Cal / 1000 mL (1.1 QT) Ready-To-Hang Prefilled Container / 8 ct
65007	Pivot 1.5 Cal / 8-fl-oz (237-mL) Recloseable Carton / 24 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1000 mL (1.1 QT)	
	Value	%RDI / %DV	Value	%RDI / %DV
Protein, g	22.2		93.8	
Fat, g	12.0		50.8	
Carbohydrate, g	40.9		172.4	
Dietary Fiber, g	1.8*		7.5*	
L-Carnitine, mg	36		150	
Taurine, mg	36		150	
Water, g	180		759	
Calories	355		1500	
Vitamin A, IU	2370 [†]		10000 [†]	
Beta-Carotene, mg	1.2		4.8	
Vitamin D, IU	95		400	
Vitamin E, IU	60		250	
Vitamin K, mcg	19		80	
Vitamin C, mg	72		300	
Folic Acid, mcg	145		600	
Thiamin (Vitamin B1), mg	0.54		2.3	
Riboflavin (Vitamin B2), mg	0.61		2.6	
Vitamin B6, mg	0.71		3.0	
Vitamin B12, mcg	2.2		9.0	
Niacin, mg	7.1		30	
Choline, mg	145		600	
Biotin, mcg	110		450	
Pantothenic Acid, mg	3.6		15	
Sodium, mg	330		1400	
Sodium, mEq	14.3		60.9	
Potassium, mg	475		2000	
Potassium, mEq	12.2		51.3	
Chloride, mg	380		1600	
Chloride, mEq	10.7		45.2	
Calcium, mg	240		1000	
Phosphorus, mg	240		1000	
Magnesium, mg	95		400	
Iodine, mcg	36		150	
Manganese, mg	1.2		5.0	
Copper, mg	0.48		2.0	
Zinc, mg	6.0		25	
Iron, mg	4.3		18	
Selenium, mcg	17		70	
Chromium, mcg	29		120	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1000 mL (1.1 QT)	
	Value	%RDI / %DV	Value	%RDI / %DV
Molybdenum, mcg	36		150	

Liquid Unflavored Footnotes & References

Per 8 fl oz (237 mL)

*1.8 g of fructooligosaccharides.

†1590 IU of vitamin A activity supplied by 1.2 mg of beta-carotene.

Per 1000 mL (1.1 QT)

*7.5 g of fructooligosaccharides.

†6360 IU of vitamin A activity supplied by 4.8 mg of beta-carotene.

Preparation

Instructions for Use:

8 fl oz recloseable carton

- Store unopened at room temperature; avoid extreme temperatures.
- Shake well prior to opening.
- Once opened, reclose or cover, refrigerate and use within 48 hours..

Oral Feeding:

- May be used for total or supplemental nutrition.
- May be fed at room temperature or chilled.

Tube Feeding:

- Follow physician's instructions.
- Adjust flow rate and volume according to patient's condition and tolerance.
- Feed at room temperature using a feeding pump or syringe.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- Avoid contamination during preparation and use.

Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the Instructions for Use increases the potential for microbial contamination and reduces hangtime.

- Administer product at room temperature.
- THOROUGHLY wash hands with soap and water before handling container or feeding set.
- Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps:

- Remove dust cover from RTH Safety Screw Cap.
- Remove dust cover from feeding set connector.
- Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- Turn connector collar clockwise until it is securely fastened.
- Close clamp on set before inverting container.
- Invert container and suspend, using hanging ring on bottom of container.
- Follow directions for use provided by manufacturer of feeding sets.
- Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.