

Product Information: Pulmocare®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

- 1 PULMOCARE is designed for people with chronic obstructive pulmonary disease (COPD), cystic fibrosis, or respiratory failure who may benefit from a high-calorie, modified carbohydrate and fat enteral formula that may help reduce diet-induced carbon dioxide production.
- 1 For tube or oral feeding.
- 1 For supplemental or sole-source nutrition.
- 1 Use under medical supervision.
- 1 Provides complete and balanced nutrition.
- 1 Meets or exceeds 100% of the 1995 RDIs for 24 essential vitamins and minerals in 1420 Cal (947 mL).
- 1 20% of fat as MCTs to enhance fat absorption.
- 1 Elevated antioxidants vitamins C & E, and beta-carotene.
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not for IV use.
- 1 Not for people with galactosemia.

Ingredients

Liquid Vanilla:

Water, Sodium & Calcium Caseinates, Sugar, Canola Oil, Corn Maltodextrin, Medium-Chain Triglycerides, Corn Oil, High Oleic Safflower Oil, Natural & Artificial Flavor, Magnesium Chloride, Calcium Phosphate, Soy Lecithin, Potassium Citrate, Sodium Citrate, Potassium Phosphate, Sodium Hydroxide, Ascorbic Acid, Choline Chloride, Potassium Hydroxide, Taurine, L-Carnitine, Gellan Gum, Salt, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Beta-Carotene, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin B12, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
62725	Pulmocare / 1000 mL (1.1 QT) Ready-To-Hang Prefilled Container / 8 ct
64811	Pulmocare Vanilla / 8 fl oz (237 mL) Recloseable Carton / 24 ct

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)	
	Value	%RDI / %DV
Protein, g	14.8	
Fat, g	22.1	
Carbohydrate, g	25.0	
L-Carnitine, mg	36	
Taurine, mg	36	
Water, g	186	
Calories	355	
Vitamin A, IU	2840*	
Beta-Carotene, mg	1.2	
Vitamin D, IU	100	
Vitamin E, IU	20	
Vitamin K, mcg	20	
Vitamin C, mg	75	
Folic Acid, mcg	200	
Thiamin (Vitamin B1), mg	0.75	
Riboflavin (Vitamin B2), mg	0.85	
Vitamin B6, mg	1.0	
Vitamin B12, mcg	3.0	
Niacin, mg	10	
Choline, mg	150	
Biotin, mcg	150	
Pantothenic Acid, mg	5.0	
Sodium, mg	310	
Sodium, mEq	13.5	
Potassium, mg	465	
Potassium, mEq	11.9	
Chloride, mg	400	
Chloride, mEq	11.3	
Calcium, mg	250	
Phosphorus, mg	250	
Magnesium, mg	100	
Iodine, mcg	38	
Manganese, mg	1.3	
Copper, mg	0.50	
Zinc, mg	5.7	
Iron, mg	4.5	
Selenium, mcg	18	
Chromium, mcg	30	
Molybdenum, mcg	38	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Liquid Vanilla Footnotes & References

Per 8 fl oz (237 mL)

*1590 IU of vitamin A activity supplied by 1.2 mg of beta-carotene.

Preparation

Instructions for Use:

8 fl oz recloseable container

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Shake well prior to opening.
- 1 Once opened, reclose or cover, refrigerate and **use within 48 hours**.

Oral Feeding:

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Feed at room temperature using a feeding pump or syringe.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.

Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the Instructions for Use increases the potential for microbial contamination and reduces hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from RTH Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.
- 1 Follow directions for use provided by manufacturer of feeding sets.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.