



Search Mayo Clinic

MENU

Log in to Patient Account

Request an Appointment

Find a Doctor

Find a Job

Give Now

English

[Patient Care & Health Information](#) [Diseases & Conditions](#)

# Urinary incontinence

[Request an Appointment](#)

[Symptoms & causes](#)

[Diagnosis & treatment](#)

[Doctors & departments](#)

[Care at Mayo Clinic](#)

## Overview

[Print](#)

Advertisement

Urinary incontinence — the loss of bladder control — is a common and often embarrassing problem. The severity ranges from occasionally leaking urine when you cough or sneeze to having an urge to urinate that's so sudden and strong you don't get to a toilet in time.

Though it occurs more often as people get older, urinary incontinence isn't an inevitable consequence of aging. If urinary incontinence affects your daily activities, don't hesitate to see your doctor. For most people, simple lifestyle changes or medical treatment can ease discomfort or stop urinary incontinence.

[Urinary incontinence care at Mayo Clinic.](#)

### Products & Services

[Book: Mayo Clinic on Managing Incontinence](#)

[Show more products from Mayo Clinic](#)

## Symptoms

Many people experience occasional, minor leaks of urine. Others may lose small to moderate amounts of urine more frequently.

Types of urinary incontinence include:

- **Stress incontinence.** Urine leaks when you exert pressure on your bladder by coughing, sneezing, laughing, exercising or lifting something heavy.
- **Urge incontinence.** You have a sudden, intense urge to urinate followed by an involuntary loss of urine. You may need to urinate often, including throughout the night. Urge incontinence may be caused by a minor condition, such as infection, or a more-severe condition such as a neurologic disorder or diabetes.
- **Overflow incontinence.** You experience frequent or constant dribbling of urine due to a bladder that doesn't empty completely.
- **Functional incontinence.** A physical or mental impairment keeps you from making it to the toilet in time. For example, if you have severe arthritis, you may not be able to unbutton your pants quickly enough.
- **Mixed incontinence.** You experience more than one type of urinary incontinence.

## When to see a doctor

You may feel uncomfortable discussing incontinence with your doctor. But if incontinence is frequent or is affecting your quality of life, it's important to seek medical advice because urinary incontinence may:

- Indicate a more-serious underlying condition
- Cause you to restrict your activities and limit your social interactions
- Increase the risk of falls in older adults as they rush to the toilet

## [Request an Appointment at Mayo Clinic](#)

## Causes

Urinary incontinence isn't a disease, it's a symptom. It can be caused by everyday habits, underlying medical conditions or physical problems. A thorough evaluation by your doctor can help determine what's behind your incontinence.

Mayo Clinic does not endorse companies or products. Advertising revenue supports our not-for-profit mission.

### Advertising & Sponsorship

[Policy](#) | [Opportunities](#) | [Ad Choices](#)

## Mayo Clinic Marketplace

Check out these best-sellers and special offers on books and newsletters from Mayo Clinic.

[The Mayo Clinic Diet Online](#)

---

[4 simple steps to a joy-filled life](#)

---

[Conquer heart disease in 10 min. a day](#)

---

[NEW – Guide to Fibromyalgia](#)

---

[Stop osteoporosis in its tracks](#)

## Temporary urinary incontinence

Certain drinks, foods and medications may act as diuretics — stimulating your bladder and increasing your volume of urine. They include:

- Alcohol
- Caffeine
- Carbonated drinks and sparkling water
- Artificial sweeteners
- Chocolate
- Chili peppers
- Foods that are high in spice, sugar or acid, especially citrus fruits
- Heart and blood pressure medications, sedatives, and muscle relaxants
- Large doses of vitamin C


Urinary incontinence may also be caused by an easily treatable medical condition, such as:

- **Urinary tract infection.** Infections can irritate your bladder, causing you to have strong urges to urinate, and sometimes incontinence.
- **Constipation.** The rectum is located near the bladder and shares many of the same nerves. Hard, compacted stool in your rectum causes these nerves to be overactive and increase urinary frequency.

## Persistent urinary incontinence

Urinary incontinence can also be a persistent condition caused by underlying physical problems or changes, including:

- **Pregnancy.** Hormonal changes and the increased weight of the fetus can lead to stress incontinence.




© 2010 FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH. ALL RIGHTS RESERVED.

### Female urinary system

Your urinary system — which includes the kidneys, ureters, bladder and urethra — is responsible for removing waste from your body through urine. Your kidneys, located toward the back in your upper abdomen, produce urine by filtering waste and fluid from your blood. That urine then travels through your ureters to your bladder, where the urine is stored until you can eliminate it at an appropriate time.



AdChc



© 2010 FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH. ALL RIGHTS RESERVED.

### Male urinary system

Your urinary system — which includes the kidneys, ureters, bladder and urethra — is responsible for removing waste from your body through urine. Your kidneys, located toward the back in your upper abdomen, produce urine by filtering waste and fluid from your blood. That urine then travels through your ureters to your bladder, where the urine is stored until you can eliminate it at an appropriate time.

- **Childbirth.** Vaginal delivery can weaken muscles needed for bladder control and also damage bladder nerves and supportive tissue, leading to a dropped (prolapsed) pelvic floor. With prolapse, the bladder, uterus, rectum or small intestine can get pushed down from the usual position and protrude into the vagina. Such protrusions can be associated with incontinence.
- **Changes with age.** Aging of the bladder muscle can decrease the bladder's capacity to store urine. Also, involuntary bladder contractions become more frequent as you get older.
- **Menopause.** After menopause women produce less estrogen, a hormone that helps keep the lining of the bladder and urethra healthy. Deterioration of these tissues can aggravate incontinence.
- **Hysterectomy.** In women, the bladder and uterus are supported by many of the same muscles and ligaments. Any surgery that involves a woman's reproductive system, including removal of the uterus, may damage the supporting pelvic floor muscles, which can lead to incontinence.
- **Enlarged prostate.** Especially in older men, incontinence often stems from enlargement of the prostate gland, a condition known as benign prostatic hyperplasia.
- **Prostate cancer.** In men, stress incontinence or urge incontinence can be associated with untreated prostate cancer. But more often, incontinence is a side effect of treatments for prostate cancer.
- **Obstruction.** A tumor anywhere along your urinary tract can block the normal flow of urine, leading to overflow incontinence. Urinary stones — hard, stone-like masses that form in the bladder — sometimes cause urine leakage.
- **Neurological disorders.** Multiple sclerosis, Parkinson's disease, a stroke, a brain tumor or a spinal injury can interfere with nerve signals involved in bladder control, causing urinary incontinence.

## Risk factors

Factors that increase your risk of developing urinary incontinence include:

- **Gender.** Women are more likely to have stress incontinence. Pregnancy, childbirth, menopause and normal female anatomy account for this difference. However, men with prostate gland problems are at increased risk of urge and overflow incontinence.
- **Age.** As you get older, the muscles in your bladder and urethra lose some of their strength. Changes with age reduce how much your bladder can hold and increase the chances of involuntary urine release.

- **Being overweight.** Extra weight increases pressure on your bladder and surrounding muscles, which weakens them and allows urine to leak out when you cough or sneeze.
- **Smoking.** Tobacco use may increase your risk of urinary incontinence.
- **Family history.** If a close family member has urinary incontinence, especially urge incontinence, your risk of developing the condition is higher.
- **Other diseases.** Neurological disease or diabetes may increase your risk of incontinence.

## Complications

Complications of chronic urinary incontinence include:

- **Skin problems.** Rashes, skin infections and sores can develop from constantly wet skin.
- **Urinary tract infections.** Incontinence increases your risk of repeated urinary tract infections.
- **Impacts on your personal life.** Urinary incontinence can affect your social, work and personal relationships.

## Prevention

Urinary incontinence isn't always preventable. However, to help decrease your risk:

- Maintain a healthy weight
- Practice pelvic floor exercises
- Avoid bladder irritants, such as caffeine, alcohol and acidic foods
- Eat more fiber, which can prevent constipation, a cause of urinary incontinence
- Don't smoke, or seek help to quit smoking

By Mayo Clinic Staff

[Urinary incontinence care at Mayo Clinic](#)

[Request an Appointment at Mayo Clinic](#)

## Diagnosis & treatment

Share on: [Facebook](#) [Twitter](#) [Print](#) April 13, 2019

[Show references](#) ▾

## Related

[Adult bed-wetting: A concern?](#)

[Bladder control problems in women: Seek treatment](#)

[Bladder control problems: Medications](#)

[Show more related content](#)

### Associated Procedures

[Cystoscopy](#)

[Ultrasound](#)

[Urinalysis](#)

### Products & Services

[Book: Mayo Clinic on Managing Incontinence](#)

[Show more products and services from Mayo Clinic](#)

**Mayo Clinic in Rochester, Minn., has been recognized as the top Urology hospital in the nation for 2019-2020 by U.S. News & World Report.**

[Learn more about this top honor](#)

---

## Urinary incontinence

---

[Symptoms & causes](#)

---

[Diagnosis & treatment](#)

Care at Mayo Clinic

[Patient Care & Health Information](#) [Diseases & Conditions](#) [Urinary incontinence](#)

CON-20305548



[Request Appointment](#) | [Contact Us](#)  
[About Mayo Clinic](#) | [Employees](#) | [Find a Job](#)  
[Site Map](#) | [About This Site](#)

Mayo Clinic is a not-for-profit organization. Make a donation.

Any use of this site constitutes your agreement to the [Terms and Conditions](#) and [Privacy Policy](#) linked below.

[Terms and Conditions](#)

[Privacy Policy](#)

[Notice of Privacy Practices](#)

[Notice of Nondiscrimination](#)

A single copy of these materials may be reprinted for noncommercial personal use only. "Mayo," "Mayo Clinic," "MayoClinic.org," "Mayo Clinic Healthy Living," and the triple-shield Mayo Clinic logo are trademarks of Mayo Foundation for Medical Education and Research.



This site complies with the [HONcode standard for trustworthy health information](#): [verify here.](#)