

Genuine Bayer[®] Aspirin Tablets

Drug Facts
Active ingredient (in each tablet) Purposes Aspirin 325 mg (NSAID)* Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug
Uses • temporarily relieves minor aches and pains due to: • headache • menstrual pain • toothache • minor pain of arthritis • temporarily reduces fever
WarningsReye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.Allergy alert: Aspirin may cause a severe allergic reaction which may include:• hives • facial swelling • asthma (wheezing) • shock
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed
 Do not use if you are allergic to aspirin or any other pain reliever/fever reducer if you have ever had an allergic reaction to this product or any of its ingredients
 Ask a doctor before use if stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic you have asthma
Ask a doctor or pharmacist before use if you are taking a prescription drug for ● gout ● diabetes ● arthritis
 Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur ringing in the ears or a loss of hearing occurs



Genuine Bayer[®] Aspirin Tablets

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
 Directions drink a full glass of water with each dose adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours children under 12 years: consult a doctor
Other information save carton for full directions and warnings Store at room temperature
<i>Inactive ingredients</i> corn starch, hypromellose, powdered cellulose, triacetin
Questions or comments? 1-800-331-4536 (Mon – Fri 9AM – 5PM EST)