

The key to safer adhesive removal

- Quick
- Painless
- Easy



detachol[®]
ADHESIVE REMOVER



UNLOCK THE SOLUTION TO HIDDEN THREATS

“Detachol... has decreased skin tears and increased patient satisfaction.”

Lori B
Pediatric CNS
Madison, WI



Detachol helps remove leftover adhesive residue, which can harbor microorganisms.

By effectively removing it, Detachol can help reduce the risk of infection.

Trauma resulting from adhesive tape and dressing removal can

- Increase the size of wounds
- Exacerbate wound pain
- Delay healing¹

Additional skin threats

- Adhesive tape has long been recognized as a potential source of nosocomial infections^{2,3}
- Stripping cells from the epidermis can cause skin tears⁴
 - When skin is exposed to moisture, the risk of bacterial colonization increases⁵
 - Stripping on vulnerable periwound skin can lead to inflammatory skin damage, edema, soreness, and pain^{4,5}

Dressing changes are often painful⁶

- Patient pain from skin trauma and tears from adhesive tape removal can be avoided
- Use of a solvent to break the bond between adhesive and skin is recommended as an approach to prevent injury^{5,7,8}



detachol
ADHESIVE REMOVER

DETACHOL OPENS THE WAY TO OPTIMAL PATIENT COMFORT AND OUTCOMES

“{Detachol}... left no residue on the skin, and allowed for much more time efficient care...”

Diane P
RN, WOCN
Aurora, CO

Maximize comfort and ease of use Minimize risk of infection

- Quickly, effortlessly, and completely removes adhesive and its residue
- Prevents skin tears that contribute to the risk of local infection; won't compromise skin barrier
- Completely removes residue in which bacteria can breed
- Reduces patient pain and discomfort

Balance patient protection and resource preservation

- Can reduce physical, emotional, and monetary costs of adhesive removal-related, hospital-acquired skin infection
- Is pain-free for pediatric patients, geriatric patients, and sensitive skin types
- Does not irritate or dry the skin⁹
- Quickly and easily removes adhesive to save healthcare professional time
- No harsh alcohol or acetone
- Will not degrade sterile gloves like alcohol-based adhesive removers

Use Detachol for safe and painless adhesive removal



Central line dressings



NPWT dressings



Catheter securement devices



Nasogastric tube securements



Peripheral IV dressings



Endotracheal tube securements
And many more!

DETACHOL: THE KEY TO SUCCESSFUL ADHESIVE REMOVAL



- Completely removes adhesive residue
- Reduces the chance of skin tears and infections
- Saves time and reduces patient pain and anxiety
- Is safe for all patient types including geriatric and pediatric patients

Available in several convenient configurations



ORDERING INFORMATION

Description	Unit	HRI #
Detachol Vials	48 X 2/3 mL	0496-0513-48
Detachol Unit Dose	15mL Bottle with plug	0496-0513-15
Detachol Dispenser Cap	2oz Bottle	0496-0513-06
Detachol Dispenser Cap	4oz Bottle	0496-0513-04

Contact Eloquest Healthcare® toll free: 1.877.433.7626, ext. 298 for product inquiries, ext. 234 for customer service OR www.detachol.com

detachol[®]
ADHESIVE REMOVER

Eloquest
HEALTHCARE[®]

Our product portfolio includes:

- Mastisol[®] Liquid Adhesive
- Detachol[®] Adhesive Remover
- Microcyn[®] Rx Skin and Wound Cleanser
- LMX4[®] Topical Anesthetic

www.eloquesthealthcare.com

REFERENCES

1. Rippon M, White R, Davies P. *Wounds UK*. 2007;3(4):76-86.
2. Berkowitz DM, Lee WS, Pazin GJ, Yee RB, Ho M. *Appl Microbiol*. 1974;28(4):651-654.
3. Redelmeier DA, Livesley NJ. *J Gen Intern Med*. 1999;14(6):373-375.
4. Lawton S, Langoen A. Assessing and managing vulnerable periwound skin. *World Wide Wounds*. October 2009. Accessed October 3, 2012. Available at: <http://www.worldwidewounds.com/2009/October/Lawton-Langoen/vulnerable-skin-2.html>.
5. Bryant RA. Types of skin damage and differential diagnosis. In: Bryan RA, Nix DP, eds. *Acute and Chronic Wounds: Current Management Concepts*. St. Louis, MO: Elsevier Mosby; 2012: chap 5:83-107.
6. Coha T, Bryant RA, Nix DP. Skin care needs of the pediatric and neonatal patient. Part I: The pediatric patient. In: Bryan RA, Nix DP, eds. *Acute and Chronic Wounds: Current Management Concepts*. St. Louis MO: Elsevier Mosby; 2012:chap 36:485-502.
7. Pennsylvania Patient Safety Authority, Patient Safety Advisory. Skin tears: the clinical challenge. Accessed October 3, 2012. Available at: [http://patientsafetyauthority.org/ADVISORIES/AdvisoryLibrary/2006/Sep3\(3\)/Pages/01b.aspx](http://patientsafetyauthority.org/ADVISORIES/AdvisoryLibrary/2006/Sep3(3)/Pages/01b.aspx).
8. Rolstad BS, Bryant RA, Nix DP. Topical management. In: Bryan RA, Nix DP, eds. *Acute and Chronic Wounds: Current Management Concepts*. St. Louis, MO: Elsevier Mosby; 2012: chap 18:289-306.
9. Fowler JF. Data on file. Eloquest Healthcare, Inc; 2011.



Scan here to watch the Detachol instructional video or visit www.detachol.com to see for yourself

Detachol[®], Mastisol[®], and LXM4[®] are registered trademarks of Ferndale IP, Inc. Microcyn[®] is a registered trademark of Oculus Innovative Sciences, Inc.

© 2013 Eloquest Healthcare[®], Inc. DE16 R06/13