

3M™ Steri-Strip™ Antimicrobial Skin Closures (reinforced)

Commonly Asked Questions

- **Components**

Question:

Do Steri-Strip Antimicrobial skin closures contain latex?

Answer:

No. Neither the packaging nor the Steri-Strip Antimicrobial skin closures contain natural rubber latex.

Question:

Do Steri-Strip Antimicrobial skin closures contain fiberglass?

Answer:

No. Steri-Strips skin closures have never contained fiberglass. The reinforcing filaments in the reinforced Steri-Strip skin closures and Steri-Strip Antimicrobial skin closures are made of polyester.

Question:

Are Steri-Strip Antimicrobial skin closures hypoallergenic?

Answer:

No. The Steri-Strip Skin Antimicrobial skin closures contain an iodophor compound, so they are not recommended for use on patients with known sensitivity to iodine.

Question:

What is the shelf life of Steri-Strip Antimicrobial skin closures?

Answer:

2 years. The expiration date may be found on the top of each Steri-Strip skin closure package. Without breaking the seal, fold the rear package flap back. Holding the package up to the light, you will note:

LOT, hourglass, year, 1-2 numbers, several letters

The hourglass symbol represents the expiration date. 2004-10 would indicate that the product will expire in October of 2004.

Question:

Are Steri-Strip Antimicrobial skin closures sterile? Can Steri-Strip Antimicrobial skin closures be resterilized?

Answer:

Steri-Strip Antimicrobial skin closures are sold sterile. They may NOT be resterilized with ethylene oxide.

- **Selection and application**

Question:

What is the best way to get great adhesion of Steri-Strip skin closures?

Answer:

Steri-Strip skin closures adhere best to clean, dry skin. This can be accomplished by ensuring that bleeding is under control and cleaning the skin with saline or alcohol to remove oils or exudates followed by gentle drying of the skin surface.

Steri-Strip closures have a “pressure sensitive” acrylate adhesive. Adhesion of skin closures will be enhanced if you gently but firmly press or stroke the skin closure strip during application so that the adhesive gets into the “nooks and crannies” of the skin surface. See application instructions. Steri-Strip Antimicrobial skin closures have the highest adhesion of the Steri-Strip skin closure product line.

Steri-Strip skin closures were designed for closure of low-tension wounds. They are contraindicated for use in high-tension wounds that cannot easily be approximated with fingers or forceps or on infected wounds.

Question:

Should I use a skin tackifier under Steri-Strip skin closures?

Answer:

Compound benzoin tincture (CBT) may be used to increase adhesion of Steri-Strip skin closures. Although, CBT does not contain iodine, it may be an irritant to some individuals and should not enter the wound.

Question:

What can I do to protect very fragile skin?

Answer:

Apply the skin closure to dry, clean skin that is free of chemicals. If hair is to be removed, a clipper should be used. To further protect from adhesive trauma, a skin barrier such as 3M™ Cavilon™ No Sting Barrier Film may be applied and allowed to dry thoroughly before applying the skin closure. Avoid chemicals designed to increase adhesion. To reduce the risk of skin stripping, gently remove Steri-Strip skin closures: “low and slow”, towards the center of the wound, while supporting the skin.

Question:

What could cause blistering of the skin under the ends of skin closure strips?

Answer:

The most common cause of blistering is tension. Edema, hematoma, or distention due to bloating may distort the skin surface. If the adhesive is firmly attached to the epidermis and the adhesive closure backing does not “give”, the epidermis may be pulled away from the dermis resulting in blisters or skin tears. Blistering is usually noted on both ends of the strip. A similar situation occurs when a joint or other area of movement is covered with a strip that doesn’t stretch. If tension is noted, loosen, reposition, or replace the skin closure strips. To decrease the risk of tension injuries:

- Do not use Steri-Strip skin closures on high-tension wounds.
- Apply Steri-Strip skin closures without tension. Do not strap the wound closed.
- Consider 3M™ Steri-Strip™ Elastic Skin Closures if edema, hematoma, distention, or movement are anticipated.
- Monitor the wound for evidence of swelling or blistering.
- If tension is noted under Steri-Strip closures, they may be replaced under the direction of a health care professional.

Question:

I suspect that the area around the wound may swell. Will this affect the type of skin closure I should use?

Answer:

Yes, if expansion or movement (such as over a joint) is anticipated, consider using Steri-Strip Elastic skin closures.

- **Length of wear**

Question:

How long can Steri-Strip Antimicrobial skin closures be worn?

Answer:

Steri-Strip skin closures are usually worn until they fall off or the healthcare provider removes them. This usually occurs within 7 to 14 days for the Steri-Strip Antimicrobial closures. Wear time may vary depending on area of the body, skin type, degree of friction to that area, etc.

Some health care providers prefer to remove, and sometimes replace, the Steri-Strip skin closures, on follow-up visits. The timing of strip removal and the length of continued reinforcement and support of the wound with skin closure strips is left to the discretion of the health care provider.

Question:

If skin closure ends loosen, should they be trimmed?

Answer:

Yes. Once the edges loosen, a skin closure strip is more likely come off than if the strip is intact. If the loosened ends of the strips are trimmed so that they do not catch on clothing, the duration of adherence is extended. However, when less than 1/2 inch of the strip remains on either side of the wound, health care providers may want to consider gently removing the strip and replacing it with a new strip.

Question:

Can I get Steri-Strip skin closures wet?

Answer:

Unless contraindicated, you may shower 24 hours after the Steri-Strip skin closures have been applied. Then pat the strips dry. However, some studies indicate that strip adherence may be improved when bathing is limited. For additional protection, you may cover the Steri-Strip skin closure with a 3M™ Tegaderm™ Transparent Dressing.

[Click here to view detailed product descriptions and prices.](#)

Call 800-397-5899 to order today!



Health Care

3M Center, Building 275-4W-02
St. Paul, MN 55144-1000
USA
1 800 228-3957
www.3M.com/healthcare

For More Information

Contact your 3M Health Care Sales Representative, or call the 3M Health Care Customer Helpline at **1-800-228-3957**. These products can be ordered from your local distributor. Outside the United States, contact the local 3M subsidiary.